



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

329. Proposed by JOHN JAMES QUINN, Ph. D., Scottsdale, Pa.

1. Determine the equation of the locus of a fixed point in a circle of radius r , rolling along the axis of an upright cylinder of the same radius, while the axis revolves (carrying the circle with it) through an angle equal to the central angle of the rolling circle formed by the radii to the fixed point and the point of contact.

2. Suppose the point projected into the surface of the cylinder.

3. What is the surface generated by the radius of the rolling circle?

4. What is the surface generated by a radius of the cylinder through the moving point?

Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

Let a = the distance between the axis of the cylinder and the line about which it revolves. Also suppose the plane of the circle is determined by the line a and the axis of the cylinder. Let the circle, in its initial position, be in the xz plane with the point of contact in the same plane.

The position of the moving point at any time, using the axis of the cylinder as initial line and initial point of contact as origin, is $r\theta - r\sin\theta$, and $r(1 - \cos\theta)$. We use the double sign, plus when the circle is not between the z -axis and the axis of the cylinder, and the minus when it is.

1. The locus is determined by $x = [a \pm r(1 - \cos\theta)]\cos\theta$, $y = [a \pm r(1 - \cos\theta)]\sin\theta$, $z = r\theta - r\sin\theta$, θ = central angle of circle.

2. If point is projected in surface of cylinder in plane of circle, the locus is determined by $x = (a \pm r)\cos\theta$, $y = (a \pm r)\sin\theta$, $z = r(\theta - \sin\theta)$.

3. Coordinates of the center of gravity of the moving radius of the circle are, $x = [a \pm r(1 - \frac{1}{2}\cos\theta)]\cos\theta$, $y = [a \pm r(1 - \frac{1}{2}\cos\theta)]\sin\theta$, $z = r(\theta - \frac{1}{2}\sin\theta)$.

\therefore Surface = rS where S is given as follows:

$$S = \int_0^\theta [4a^2 \pm 8ar + 8r^3 \mp 4ar\cos\theta(4\sin^2\theta - 1) + r^2(1 - 16\cos\theta)(1 + \cos^2\theta)]^{\frac{1}{2}} d\theta.$$

4. Coordinates of center of gravity of radius of cylinder through moving point are $x = (a \pm \frac{1}{2}r)\cos\theta$, $y = (a \pm \frac{1}{2}r)\sin\theta$, $z = r(\theta - \sin\theta)$.

\therefore Surface = rS where

$$S = \int_0^\theta [(a \pm \frac{1}{2}r)^2 + r^2(1 - \cos\theta)^2]^{\frac{1}{2}} d\theta.$$

Also solved by H. V. Spunar.

CALCULUS.

257. Proposed by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

If $A = \int_0^\infty \frac{dx}{\sqrt{x}(2a+x)^n}$, $B = \int_0^\infty \frac{y^n dy}{\sqrt{y}(a^2+y^2)^n}$, find A/B .

Solution by J. SCHEFFER, A. M., Hagerstown, Md., and the PROPOSER.

Let $x=2a\tan^2\theta$, $y=a\tan\phi$, $m=\frac{1}{2}(2n-1)$.

$$\therefore A = \frac{2}{(2a)^{n-\frac{1}{2}}} \int_0^{\frac{1}{2}\pi} \cos^{2(n-1)}\theta \, d\theta = \frac{1}{2(2a)^m} \int_0^{\frac{1}{2}\pi} \cos^{2m-1}\theta \, d\theta = \frac{\sqrt{\pi} \Gamma(m)}{(2a)^m \Gamma\left(\frac{2m+1}{2}\right)}.$$

$$\begin{aligned} B &= \frac{1}{a^{n-\frac{1}{2}}} \int_0^{\frac{1}{2}\pi} \tan^{n-\frac{1}{2}}\phi \cos^{2(n-1)}\phi \, d\phi = \frac{1}{a^m} \int_0^{\frac{1}{2}\pi} \sin^m\phi \cos^{m-1}\phi \, d\phi \\ &= \frac{\Gamma\left(\frac{m+1}{2}\right) \Gamma\left(\frac{m}{2}\right)}{2a^m \Gamma\left(\frac{2m+1}{2}\right)}. \end{aligned}$$

$$\text{But } \Gamma\left(\frac{m+1}{2}\right) \Gamma\left(\frac{m}{2}\right) = \frac{\sqrt{\pi} \Gamma(m)}{2^{m-1}}.$$

$$\therefore B = \frac{\sqrt{\pi} \Gamma(m)}{(2a)^m \Gamma\left(\frac{2m+1}{2}\right)} = A. \quad \text{Hence } A/B=1.$$

Exhaustive solutions, though differing in results from the one given here, were received from M. V. Spunar, Francis Rust, and T. G. Wodo.

258. Proposed by A. H. HOLMES, Brunswick, Maine.

$$\text{Evaluate } \int_0^{\frac{1}{2}\pi} \frac{dx}{\sqrt{[2ax-x^2 \sqrt{a^2-x^2}]}}.$$

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Let $x=a\sin\theta$. Then

$$\begin{aligned} I &= \int_0^{\frac{1}{2}\pi} \frac{dx}{\sqrt{[2ax-x^2 \sqrt{a^2-x^2}]}} = \int_0^{\frac{1}{2}\pi} \frac{\cos\theta \, d\theta}{\sqrt{[2a\sin\theta - a\sin^2\theta \cos\theta]}} \\ \therefore I &= \frac{1}{\sqrt{2}} \int_0^{\frac{1}{2}\pi} \left[1 + \frac{a}{4}\sin\theta \cos\theta + \frac{3a^2}{32}\sin^2\theta \cos^2\theta + \frac{5a^3}{128}\sin^3\theta \cos^3\theta \right. \\ &\quad \left. + \frac{35a^4}{2048}\sin^4\theta \cos^4\theta + \frac{64a^5}{8192}\sin^5\theta \cos^5\theta + \frac{231a^6}{65536}\sin^6\theta \cos^6\theta + \dots \right] \frac{\cos\theta \, d\theta}{\sqrt{(\sin\theta)}} \\ &= 1 + \frac{31}{15} \cdot \frac{a^2}{2^9} + \frac{85015}{1989} \cdot \frac{a^4}{2^{19}} + \frac{2350494}{38675} \cdot \frac{a^6}{2^{28}} + \dots + \frac{1}{\sqrt{2}} \int_0^{\frac{1}{2}\pi} \left[\frac{a}{4}\sin\theta \cos\theta \right. \end{aligned}$$